Microfluidics for chemical and biological engineering



Shepherd, Conrad, Lewis, *et al.*, *Langmuir* **22**, 8618–8622 (**2006**)

Jacinta C. Conrad

University of Houston
Chemical and Biomolecular Engineering
CHEE 1131
Fall 2015

Fluid: physical definition

A fluid is a material that flows under an applied stress

Liquid: constant volume

Gas: volume of container

http://sciencekids.co.nz

Two physical properties of fluids:

- Viscosity: measure of fluid resistance to stress μ [mass/length-time]
- Density: ρ [mass/length^3]

Macroscale flows

Characteristics:

- Large length scales L
- Fast flow speeds V
- Turbulent flow

Many macroscale flows are characterized by large Reynolds number:

$$\text{Reynolds number Re} = \frac{\text{inertial force}}{\text{viscous force}} = \frac{\rho V L}{\mu} \gg 1\text{: turbulent}$$

Where do flows appear in a chemical plant?

http://www.photo-dictionary.com/photofiles/list/687/1097petrochemical_plant.jpg

Flow examples in plants (unit operations)

- Combination: mixing operation to create a homogeneous system
  - Requires control over mixing streams
- Separation: separation of mixture components
  - Emulsification: creation of a liquid-in-liquid suspension
  - Distillation: separation of one liquid from another liquid
  - Evaporation: removal of a gas from a mixture
- Reaction: reaction among chemical species in a mixture
  - Synthesis: e.g. creation of particles or chemicals

Microfluidics: miniaturization of flows

The introduction of microfluidics or lab-on-a-chip devices allows unit operations to be carried out in a small format:



plant: meters to kilometers
piping: cm to m

device: mm to cm
channels: μm to mm

“Miniaturization puts chemical plants where you want them”: R. F. Service, *Science* 202, 400 (1998)

Length scales for microfluidic flows



Nguyen and Wereley, *Fundamentals and Applications of Microfluidics*, 2nd ed. (2006)

Materials for microfluidics: elastomers



Advantages:

- Easy to prototype and replicate (via soft lithography)
- Cheap materials (polydimethylsiloxane, commercially available)

Disadvantages:

- Flexible and deformable (poor for high-pressure applications)
- Poor resistance to organic solvents

Microfluidic physics is different! 1

University of New Mexico (DeMoss and Cahill) (http://panda.unm.edu/flash/viscosity.phtml)

Critical flow properties in devices

Reynolds number Re = $\frac{\text{inertial force}}{\text{viscous force}} = \frac{\rho V L}{\mu}$ ≪ 1: laminar flow

Physical meaning: fluid elements follow straight streamlines, and fluid interfaces remain nearly parallel over long distances in microfluidic devices

Microscale flow physics is different! 2



Critical flow properties in devices

Reynolds number Re = $\dfrac{\text{inertial force}}{\text{viscous force}} = \dfrac{\rho V L}{\mu}$ « 1: laminar flow

Physical meaning: fluid elements follow straight streamlines, and fluid interfaces remain nearly parallel over long distances in microfluidic devices

Péclet number Pe = $\dfrac{\text{time to diffuse}}{\text{time to convect}} = \dfrac{V L}{D_0}$ » 1: fast convection

Physical meaning: diffusion is very slow compared to convection in microfluidic devices, and thus mixing requires special device designs

Combination: passive planar micromixer

Key idea: Modify geometry to obtain mixing via changing flow pattern



Melin *et al.*, *Lab Chip* **4**, 214-219 (2004)

Combination: parallel lamination mixer

Key idea: Split streams to increase surface area and hence mixing



Kamholz *et al.*, *Anal. Chem.* **71**, 5340-5347 (1999)



Schwesinger *et al.*, *J. Micromech. Microeng.* **6**, 99-102 (102)

Combination: 3-D microvascular networks

Key idea: Split streams in 3-d geometries to enhance mixing



Therriault *et al.*, *Nat. Mater.* **2**, 265-271 (2003)

Combination: herringbone micromixers

Key idea: Add elements to “fold” fluid via chaotic advection

Slanted ridges:



3-D herringbone:





Stroock *et al.*, *Science* **295**, 647-651 (2002)

Combination: herringbone mixer movie



Folch lab (University of Washington) via YouTube

Combination: microfluidic valving

<u>Key idea</u>: Fabricate a plastic valve that is separately actuated with air





Unger *et al.*, *Science* **288** 113-116 (2000)

Combination: colloid valves

Key idea: Incorporate micron-sized colloidal particles into devices

passive valve | actuated valve





Terray *et al.*, *Science* **296**, 1841-1844 (2002)

Combination: in-situ piston

Key idea: Photopolymerize parts in place in microfluidic devices





Hasselbrink *et al.*, *Anal. Chem.* **74** 4913-4918 (2002)

Separation: emulsification ("droplets")

Key idea: Exploit the Rayleigh-Plateau instability to create emulsion drops





Stress: elongates jet of liquid
Surface tension: minimizes surface area
Result: jet breaks up into drops

Thorsen *et al.*, *Phys. Rev. Lett.* **86** 4163-4166 (2001)

Emulsification: flow-focusing

Key idea: "Pinch off" droplets using a flow-focusing geometry



Anna *et al.*, *Anal. Chem.* **74** 4913-4918 (2002)





Abate lab (University of California San Francisco) via YouTube

Droplets + valving = adjustable sizes



Emulsification: enhanced mixing in drops

Key idea: Recirculation within drops enhances mixing rates



Tice *et al.*, *Langmuir* **19** 9127-9133 (2003)

Emulsification: drops in drops (in drops...)

<u>Key idea</u>: Encapsulate drops in other drops to create multiple emulsions



Utada *et al.*, *Science* **308** 537-541 (2005)



Chu *et al.*, *Angew. Chem. Int. Ed.* **46** 8970-8974 (2007)

Drops in drops: tune flow rates



Separation: cell sorting via optical forces

Key idea: Use radiation pressure to sort cells in a microfluidic device



Wang *et al.*, *Nat. Biotechnol.* **23** 83-87 (2005)



Separation: deterministic lateral displacement

Key idea: Particles of different diameter follow different streamlines







Huang *et al.*, *Science* **304** 987-990 (2004)

Separation of parasites from blood



Holm *et al.*, *Lab Chip* **304**, 1326-1332 (2011)

Separation: motile sperm sorter

<u>Key idea</u>: Live cells swim across laminar streamlines



Cho *et al.*, *Anal. Chem.* **75** 1671-1675 (2003)

Separation: distillation

Key idea: Establish vapor-liquid equilibrium in segmented flow and separate vapor using capillary forces



Hartman *et al.*, *Lab Chip* **9**, 1843-1849 (2009)

Reaction: drops as microreactors

Key idea: Drops increase reaction rates by increasing surface-to-volume ratio, reducing diffusion distances, and enhance heat and mass transfer







Ahmed *et al.*, *Adv. Synth. Catal.* **348** 1043-1048 (2006)

Reaction: enzyme kinetics

Key idea: Design a droplet-based microfluidic system to extract kinetic parameters of an enzymatic reaction



Song and Ismagilov, *J. Am. Chem. Soc.* **125** 14613-14619 (2003)

Reaction: nucleation

Key idea: Design a droplet-based microfluidic system to study effect of mixing on nucleation of protein crystals



Chen *et al.*, *J. Am. Chem. Soc.* **127** 9672-9673 (2005)

Reaction: nanoparticle synthesis

Key idea: Use of gas slugs to separate small liquid reaction volumes increases the monodispersity of microfluidically-produced particles



Khan *et al.*, *Langmuir* **20** 8604-8611 (2004)

Reaction: microfiber synthesis

Key idea: Photopolymerize a flow-focused stream "on the fly"





microfibers

microtubules



Khan *et al.*, *Lab Chip* **4** 576-580 (2004)

Reaction: gradient etching

Key idea: Gradients in reactant composition generate differences in etching rates through a surface



Jeon *et al.*, *Langmuir* **16** 8311-8316 (2000)

Applications of microfluidics

- Chemical synthesis
  - Especially for high-value components
- Controlled release
  - Pharmaceuticals
  - Cosmetics
- Biotechnology
  - Genomics and sequencing
  - Biodetection
  - Directed evolution
- Models of biological processes
  - Microvasculature and veination
  - Chemotaxis and chemical response

Application: crystallization

Goal of research: determine conditions and kinetic pathways for crystallization of biological proteins (e.g. xylanase)

<u>Key idea</u>: Change salt concentration “on chip” in an integrated microfluidic device to trigger crystallization





Kim *et al., J. Am. Chem. Soc.* **129**, 8825-8835 (2007)

Application: on-chip multistep synthesis

Goal of research: demonstrate optimized synthesis for sensitive compound

<u>Key idea</u>: Move all operations "on chip" in an integrated microfluidic device





- Microfluidic synthesis increased yield (38%) and purity (97.6%)
- Dramatic increase in time (14 min vs 50 min)

Lee *et al.*, *Science* **310**, 1793-1797 (2005)

Application: programmable release

Goal of research: controllably release multiple components in a pharmaceutical or cosmetic formulation

Key idea: Sequentially dissociate bilayer membranes in a double emulsion



Kim *et al.*, *J. Am. Chem. Soc.* **133** 15165-15171 (2011)

Commercialized technology: Capsum

Capsum (France) markets encapsulation technologies to luxury cosmetics manufacturers such as Amore Pacific (Korea)

capsum.eu

amorepacific.com

Application: directed evolution

Goal of research: identify mutants of horseradish peroxidase enzyme with higher catalytic activity

Key idea: Use ultrahigh throughput screening to remove inactive mutants



Agresti *et al.*, *Proc. Natl. Acad. Sci. USA* **107** 4004-4009 (2010)

- 108 enzyme reactions screened in 10 h (1,000× faster)
- Sample volume: < 150 μL of reagent (1,000,000× cheaper)

Application: cancer detection

Goal of research: capture rare circulating tumor cells (CTCs) in patients' bloodstreams for cancer detection and monitoring

<u>Key idea</u>: Increase surface encounter rate using chaotic advection



- Cancer cells detected at ~400 CTCs/mL
- Imaging-based platform identified new CTC clusters

Stott *et al.*, *Proc. Natl. Acad. Sci. USA* **107** 18392-18397 (2010)

Application: tissue engineering

Goal of research: model complex vascular phenomena, including angiogenesis and thrombosis

Key idea: Use microfluidic channels as a model for microvasculature



Zheng *et al.*, *Proc. Natl. Acad. Sci. USA* **109** 9342-9347 (2012)

Application: whole genome sequencing

Goal of research: analyze genome of single cells and microbial consortia without sample contamination

Key idea: Create multiplexed chip to sort, cultivate cells and identify, amplify, and sequence whole genomes



[46]Leung *et al.*, *Proc. Natl. Acad. Sci. USA* **109** 7665-7670 (2012)

Challenges

- Scale-up
    - Transition from “lab scale” devices to plant-scale operations
    - 2-d to 3-d layouts
- Interplay between parallelized chips
    - Need to generate uniform flow across multiple devices
    - Synchronization and chaotic effects
- Clogging and unsteady flow

Summary of lecture

- Microfluidics enables mini “chemical plants”
  - Exceptional control over reactions and mixing
  - Naturally achieves continuous production
- Optimal usages of microfluidic devices:
  - Specialty chemicals and high-value chemicals
  - Hard-to-produce molecules (especially biomolecules)
- Industries impacted by microfluidics
  - Biotechnology: genome sequencing, protein crystallization
  - Chemical synthesis: radiolabeled molecules
  - Manufacturing: designer specialty cosmetics
- Opportunities abound for chemical and biomolecular engineers to design new microfluidic processes